THE

SUMMARY PROCEEDINGS

4th ANNUAL MALARIA CONTROL PROGRAM REVIEW
Ethiopia and Nigeria

Held on March 8, 2013 at
The Carter Center, Atlanta, Georgia

OCTOBER 2013

TABLE OF CONTENTS

ACRONYMS ii

ACKNOWLEDGEMENTS iii

FRONTISPIECE PHOTOGRAPHS iv

EXECUTIVE SUMMARY 1

RECOMMENDATIONS 7

ETHIOPIA PRESENTATIONS **9-21**

Ethiopia Federal Ministry of Health Malaria Strategy and Activities 9

The Carter Center Malaria Program: Ethiopia Progress Report 13

Malaria Elimination Demonstration Project in Amhara 20

NIGERIA PRESENTATIONS **22-35**

Nigeria National Malaria Control Program Strategy and Activities 22

The Carter Center Malaria Program: Nigeria Progress Report 27

Mobilizing CDDs for Malaria Monitoring and LLIN Distribution in Plateau State 33

ANNEXES

1. Overview of Malaria Disease, the History of The Carter Center Malaria Control Program and Malaria Program Priorities 36
2. The Neglected Tropical Diseases 41
3. .....al.....

ACRONYMS

ACT	Artemisinin-based Combination Therapy
ACSM	Advocacy, Communication and Social Mobilization
AMFm	Affordable Medicine Facility – Malaria
BCC	Behavior Change Communication
CDC	Centers for Disease Control and Prevention
	Community Directed Distributors

ACKNOWLEDGEMENTS

The Carter Center's Malaria Control Program extends its sincere gratitude for the current and past support provided by partners and donors, including Chevron Corporation, Clarke Mosquito Control, the Bill & Melinda Gates Foundation, Sanofi-aventis, the Kingdom of Saudi Arabia, Starr International Foundation, Vestergaard Frandsen, Lions Clubs International Foundation, Lions Clubs of Ethiopia, and many generous individual donors. Because of the generosity of our partners, we continue to celebrate new achievements and expand our malaria control efforts each year, giving hope to millions of people in some of the most underserved communities in Ethiopia and Nigeria.

Photo 1. 2012 Carter Center malaria program review participants (March, 2013).

Photo 2. National conference to stop malaria and lymphatic filariasis in Nigeria (March, 2012).

Photo 3. Workshop for malaria elimination project in Amhara

EXECUTIVE SUMMARY

The Carter Center's (TCC) Malaria Control Program (MCP) provides support to national malaria programs in Ethiopia and Nigeria, with a particular focus on delivering and monitoring activities

many compelling arguments for the co-implementation of malaria and LF activities in Nigeria and in other African countries where the two diseases are co-endemic. Primary among these is the fact that both diseases are transmitted by the same Anopheles mosquito. Thus interventions such as LLIN distribution and other vector control activities can lead to significant reductions in the transmission of both diseases. In addition, increasing awareness that LLINs can prevent the sequalae of LF (skin problems, swollen legs, enlarged scrotum in men, and the associated stigma) is likely to result in increased LLIN use. This may be particularly the case in places like Nigeria where malaria is extremely common but the case fatality rate among adults is relatively low. Adults, especially adult males, may dread the prospect of being infected with LF much more than they fear malaria. Increased collaboration between LF and malaria programs has the potential to accelerate the scale-up of interventions for both diseases, as well as to increase programmatic cost effectiveness and cost efficiency. In March 2012, The Carter Center co-sponsored a two-day meeting, hosted by FMOH, to explore the shared opportunities between the national malaria and LF programs. The conference was chaired by former head of state General Dr. Yakubu Gowon and attended by over 200 people including Federal and State Ministry personnel, non-gov5.141 (re)-3(a)aM.-3(at) 21 Twion8laateasio0r h141 1(t)-iz

Figure 1

Ms. Solomon also presented the results of the National Malaria Indicator Survey (MIS) conducted in Ethiopia in 2011. The results indicated that gains in key malaria interventions and reductions in parasite prevalence following the scale-up of malaria control activities in 2006-2007 have been sustained. The results indicate that more than half (55.2%) of households have at least one mosquito net of any type and that nearly half (46.6%) of surveyed houses <2000m were sprayed with IRS in the past 12 months. Overall, 38.2% of children under 5 years (U5) and 35.3% of pregnant women reported sleeping under a net the previous night. Among households owning at least one net, approximately two-thirds of children U5 (64.5%) and pregnant women (64.2%) slept under a net the night before the survey. Nationwide parasite prevalence remained at around 1% in both 2007 (0.9%, 95% CI 0.5%-1.3%) and 2011 (1.3%, 95% CI 0.7%-1.8%).

Dr. Zerihun Tadesse, Country Representative for the Carter Center's Ethiopia office, summarized TCC-supported malaria control activities in Ethiopia. In 2012, the MCP in Ethiopia primarily provided support to the FMOH and the Regional Health Bureaus for malaria control activities conducted in Amhara, Oromia, Southern Nations Nationalities and People's Region (SNNPR), and Beneshangul Gumuz regions. This support consisted of routine monitoring of LLIN ownership and use, supportive supervision to improve the quality of case management and epidemic monitoring, training of health workers on malaria diagnostics and treatment and, in Amhara, mass screening and treatment of fever cases in the context of MDA campaigns for trachoma

Conclusions and Recommendations
Drs. Hopkins, Richards and Emerson

Ethiopia National Malaria Control Program

Presented by Ms. Hiwot Solomon, National Malaria Programme Focal Person, Federal Ministry of Healt
Ethiopia

Malaria is one of the leading causes of morbidity and mortality in Ethiopia. An estimated 55.7 million people (68% of the population) are at risk for malaria and approximately 80% of the 736 woredas (districts) in Ethiopia are considered "malarious". Malaria transmission is generally seasonal and unstable, though patterns and intensity of transmission vary throughout the country due to differences

vivax in non-malarious areas. First line treatment for severe malaria is intravenous quinine. Patients presenting to health posts with signs of severe malaria can be given rectal artesunate or intramuscular artemether as pre-referral treatment. In 2012, staff from 375 health centers received training on malaria diagnosis and treatment guidelines.

Table 1. Quantities of rapid diagnostic tests (RDT) and artemisinin-based combination therapy (ACT) distributed in Ethiopia, by year.

	RDTs Distributed	ACTs Distributed
2009	2,900,000	8,100,000
2010	8,000,000	10,400,000
2011	20,000,000	11,000,000
2012	17,900,000	9,000,000

Prevention. The main vector control activities implemented in Ethiopia are LLINs, IRS, and environmental control. The LLIN objectives are to ensure that 100% of households in malarious areas own at least one LLIN per sleeping space, and that at least 80% of people at risk of malaria use LLINs. Figure 2 illustrates that more than 45 million nets have been distributed in Ethiopia through 2012. Multiple mechanisms are used to ensure all families in malaria-affected areas can protect themselves from malaria including: free mass distribution of LLINs to all households in malarious areas still in need of LLINs (catch-up); replacement of old ineffective LLINs with new free LLINs through the HEP and campaigns (keep-up); free distribution of LLINs during emergencies; subsidized nets for other households, especially in urban areas through market priming/social marketing.

IRS is currently targeted to epidemic-prone areas and malaria-affected communities with low access to the health care system. In 2012, a total of 4,383,819 households (73.1% of targeted households) were sprayed with IRS—an increase over past years' performance (Figure 3). The goal was to reach 70% of targeted households by 2011 and 90% by 2013. Training is underway to facilitate decentralized IRS operation at the health post level. Insecticide resistance studies are also being conducted to provide evidence-based recommendations for

with limited and well-known breeding habitats.

Surveillance and Epidemic Control.

As shown in Table 2, nearly half (46.6%) of households <2000m were covered by indoor residual spraying (IRS) of insecticide in 2011. This was a significant increase over IRS coverage in 2007 (20.0%). However, the proportion of households <2000m that owned at least one net (of any type) declined from 68.9% in 2007 to 55.2% in 2011, while the proportion that owned more than one net (of any type) declined from 38.3% to 23.9%. These declines are attributed to deterioration of nets, which have an average lifespan of 3 years, suggesting the need for greater emphasis on net replacement (or "keep up") strategies in Ethiopia. Among all surveyed households, 38.2% of children U5 and 35.3% of pregnant women reported sleeping under a net last night. However, approximately two-thirds of children U5 (64.5%) and pregnant women (64.2%) in households <2000m owning at least one net reported sleeping under a net last night—values essentially unchanged from 2007, when 60.2% of children U5 and 64.3% of pregnant women reported sleeping under a net the previous night.

Table 2. Comparison of key results from 2007 and 2011 Malaria Indicator Surveys, Ethiopia (areas <2000m).

	2007	2011

Figure 8. Average zonal health facility reporting in Amhara region, Ethiopia.

Since implementation, there has been a 75.2% improvement in health facility reporting (Figure 8), as average health facility reporting within zones improved from 51.6% in January 2012 to 90.5% in December 2012.

Response to Surveillance Data. Analysis of ESS data prompted several malaria outbreak investigations in Amhara in 2012. While it is difficult to compare incidence from 2012 to previous years due to the significant increase in reporting and lack of historical district-level data, these alerts were driven by zone-level year-to-year comparisons and examination of district level trends from available data. TCC and ARHB investigated a rapid increase in cases across 11 districts in west Amhara beginning in May 2012. Emergency meetings were held with officials at regional, zone, and district level to form a joint plan of action. Mass fever screen, test and treat (MSAT) campaigns were conducted in 75 villages in June 2012 in accordance with national epidemic response guidelines. Out of 508,621 persons screened, 19,247 (3.8%) were identified as febrile and were tested for malaria by RDT. Of those tested, 8,367 (43.5%) were positive for Plasmodiuminfection, with 74.1% of cases due to P. falciparum. House-to-house MSAT was conducted by district health officers, with assistance from HEWs and the Health Development Army (HDA). In addition, TCC and ARHB officials visited health facilities to ensure adequate supply of RDTs and anti-malaria drugs, and directed IRS malarip(u)2(re)1(v)-(re)3(n)2(t)-31(o)-n(h)2(an)2(d)2

Table 4. Summary of MalTra Week campaigns in Amhara Region, Ethiopia, 2012.			
	MalTra Week VIII—East Amhara (May 2012)	**MalTra Week IX—West Amhara (November 2012)**	**TOTAL FOR 2012**
Number of people attending	6,193,022	7,514,787	13,707,809
Number with fever (%)	13,222 (0.2%)	77,369 (1.0%)	90,591 (0.7%)
Number RDT-positive (%			

Amhara, a total of 10,463 households were assessed in 2012. As shown in Table 5, household net

(LLIN coverage and utilization; IRS coverage) have not yet reached target thresholds, indicating the need to further support these core interventions. The project also aims to incorporate innovative approaches for malaria elimination that may include: robust surveillance systems capable of detecting clusters of cases in real-time; reactive case detection with potential MDA in areas surrounding index cases; mass screen, test, and treat activities; evaluation of ivermectin MDA on malaria incidence; and potential evaluation of novel drugs or drug formulations to target transmissible (gametocytes) and dormant (P. vivax hypnozoites) forms of the parasite in humans. The partners also recognize that entomologic

The plan also focuses on a number of cross-cutting strategies including: advocacy, communication and social mobilization (ACSM); procurement and supply chain management; monitoring and evaluation; program management; and intersectoral collaboration.

During the 2012 malaria program review meeting, the presentation by the Nigerian NMCP focused on the elements of the strategy that overlap with current Carter Center primary activities in Nigeria.

Longłasting Insecticidal Net Distribution: Scaling up for Impact (SUFI)
In

nding and procurement delac11(y)-5fs ac11(t)-3(t)-3fhe national, state andac11(L)-5fGA levels;oss of

distribute ~97 million LLINs in 2013 and 100,000,000 LLINs in 2014 through these channels, as well as through additional mass campaigns, to achieve universal coverage.

The NMCP is also implementing a plan for monitoring and managing insecticide resistance. Sentinel sites will be established for this purpose.

Case Management and Intermittent Preventive Therapy in Pregnancy (IPTp)

In 2011, Nigeria launched the first phase of the Affordable Medicines Facility-malaria (AMFm) program. AMFm enables countries to purchase first-line malaria treatments at significantly reduced costs and to pass those cost savings on to patients at both public and private sector facilities. As the result of this program, artemisinin-based combination therapies (ACTs) are now provided at no cost at some public health facilities, and the NMCP is engaged in continuous efforts to expand the number of facilities with consistent and adequate stock of RDTs and ACTs. Furthermore, the NMCP and its partner organizations are training Community-Oriented Resources Persons (CORPS) to use RDTs and to treat malaria appropriately with free ACTs based on either clinical signs or the results of a diagnostic test. The NMCP is working with Proprietary Patent Medicine Vendors to increase opportunities for the sale of efficacious malaria treatments at subsidized costs. According to malaria surveillance data from 2012, ~42% of all malaria cases captured by the surveillance system (see below) were treated with ACTs.

The NMCP is also working with partner organizations to launch a pilot study of Seasonal Malaria Chemoprophylaxis (SMC) in two states of northern Nigeria.

National Malaria Surveillance

Malaria case data in Nigeria are reported by each state through the Integrated Disease Surveillance and Response (IDSR) system. However, given the low availability of RDTs nationwide, these data primarily capture clinical malaria (fever cases) rather than confirmed malaria. Additionally, IDSR reporting by private health facilities is very poor. Given this, the NMCP has introduced a separate program-specific Roll-Back Malaria (RBM) reporting system in select health facilities. These health facilities are provided with RDTs to allow for the reporting of confirmed cases. This RBM system has been introduced in all 36 states plus the Federal Capital Territory and includes 11,610 public health facilities (~51% of all public health facilities) and 6,434 private health facilities (~5%). Some key indicators reported by these facilities on a monthly basis are listed below:

- # of children under five with fever who receive a diagnostic test
- # of children under five with confirmed uncomplicated malaria treated with ACTs
- # of children under five treated with ACTs on basis of fever only
- # of persons 5-36 years and 36+ years with fever who receive a diagnostic test
- # of persons 5-36 years and 36+ years with confirmed uncomplicated malaria treated with ACTs
- # of pregnant women receiving IPT1 & 2 (data from HMIS reports)

In 2012, this RBM surveillance system captured 13,166,487 malaria cases (confirmed and unconfirmed), which accounted for 25% of hospital attendance. Of those, 42% (n=5,579,123) were treated with ACTs. A total of 4,787 malaria-attributable deaths were reported through this system, but it is important to note that this is not a measure of the total number of deaths due to malaria in Nigeria as data are reported by only a small proportion of health facilities and many persons may die of malaria without ever seeking care from an authorized public or private facility.

Plans for improving surveillance reporting in the future include developing modalities for capturing data from a greater proportion of private facilities, engaging secondary and tertiary facilities to improve data reporting and introducing an electronic data reporting system (DHIS2).

Collaboration between Malaria and LF Programs in Nigeria

During 2012, the NMCP was actively engaged in a number of different activities designed to increase collaboration between the national malaria and lymphatic filariasis programs in Nigeria, and thus maximize potential synergies between the two programs.

The Carter Center Malaria Control Program: Nigeria Progress Report

Presentation by Mr. Adamu Sallau, Director, Malaria Control Program
The Carter Center, Nigeria

The Carter Center-Assisted Malaria Control Program in Nigeria: Priorities and Activities

The priorities of the Carter Center's (TCC) Malaria Control Program (MCP) in Nigeria are to provide assistance to the federal and state Ministries of Health for their malaria activities and to demonstrate the effectiveness of innovative approaches to malaria control in order to reduce and eventually eliminate malaria transmission in nine states assisted (v)-5(it2(in)2(n1(o)-3(ra)111.)-a)10(s)-1(t)-7(n)2(e)-3(t)8(o)-5(it21

other disease programs in malaria control activities, including LLIN distribution, monitoring and BCC. TCC's monitoring and evaluation activities include routine process evaluations as well as large-scale surveys to assess malaria and anemia prevalence, net ownership, and net use. TCC Nigeria also provides assistance with post-campaign evaluations of LLIN distributions, and provided support for a National Malaria Indicator Survey conducted in 2010. Innovative pilot projects and operational research activities conducted in Nigeria have included a study of the effectiveness of different net distribution strategies on malaria and LF transmission, a study on the cost-effectiveness of integration, and a pilot demonstration of a community-based behavior change communications intervention. The specific activities conducted in 2012 in each of the categories described above are presented in detail here.

Programmatic Interventions

LLIN Distribution

Between 2004 and 2012, TCC Nigeria provided support (technical and financial) for the distribution of a total of 7,675,963 insecticide-treated nets[7] in six states of Nigeria (Plateau, Imo, Abia, Ebonyi, Edo, and Enugu), through a combination of mass statewide campaigns conducted as part of the national scale-up and a number of smaller localized campaigns conducted as part of routine distribution or operational research projects (Figure 13). Out of the nine TCC-assisted states in Nigeria, all but one have now completed their statewide campaigns. The last remaining state, Delta, is scheduled to complete its campaign in May of 2013.



Figure 13. Annual number of insecticide treated nets (ITNs) distributed in Nigeria with support from The Carter Center between 2004 and 2012.

[7] All nets distributed since 2006 have been LLINs. However, the nets distributed prior to 2006 were insecticide-treated nets that required periodic retreatment.

Out of the cumulative total of LLINs distributed with TCC assistance, 3,319,046 were distributed in 2012, in the context of mass distribution campaigns in Imo (1,451,209 LLINs, 78.8% of state target), Abia (710,530 LLINs, 54.9% of state target) and Edo (1,157,307, 78.8% of state target). This is the greatest number of nets distributed with TCC support in a single year to date. TCC provided financial support to cover training costs, technical support for all phases of the campaigns in these states and vehicles, and also transported supplies. In addition, The Carter Center coordinated advocacy visits by former head of state General Yakubu Gow

age, compared to other age groups?; and 2) what might explain non-net-use among the subset of people with access to nets who do not use them? With regards to the first question, participants' responses suggested that net use may be lower among adolescents because they were not prioritized during net distribution and thus have limited access to nets

127 private), 318 (36%) are expected to report routine malaria data on a monthly basis through the RBM reporting system. This subset of facilities is prioritized to receive RDTs and ACTs. However, only 255 (29%) of these facilities currently receive RDTs and only 510 (7%) of the 7072 health workers in the public facilities have received training on the correct use of RDTs. (2iinn3

LLIN Monitoring During the Annual MDA Census in Plateau State

During the annual MDA census, CDDs visit all of the households in their communities to record the number, sex, and age of household members. In 2012, the census register used in Plateau state was modified to also include information on bed net ownership, number of sleeping spaces and bed net needs. In places targeted to be the first to receive a planned behavior change intervention as part of a gradual state-wide scale up of behavior change communications (BCC), CDDs also collected information on net use and care.

LLIN ownership and need data were reported for a total of 365,566 households in 3,140 (98%) of the 3,189 villages with CDDs in 15 of the 17 local government areas (LGAs) in Plateau state. Given extended health worker strikes and the challenges associated with conducting census activities in urban areas with high population density, two large and predominantly urban LGAs (Jos North and Jos South) had not yet submitted LLIN register data at the time of the program review meeting.

The LLIN register data indicate that the average household size in Plateau is 7.2 persons, that there is an average of 4.8 sleeping spaces per household and that 1.6 persons, on average, share a sleeping space. A total of 1,108,196 additional LLINs would be required to ensure that every sleeping space in the areas that reported data is covered by a net. This total does not reflect the net needs of villages without CDDs (<10% of villages according to best estimates available at the time of the review meeting) or of the two b302(<)-3(1)60.

planned nation-wide scale-up of malaria control activities, with a focus on the mass distribution of long-lasting insecticidal nets (LLINs).

A dedicated Carter Center MCP was launched in Ethiopia in February 2006. Then Ethiopian Minister of Health, Dr. Tedros Adhanom, requested that TCC join his country's national effort to provide protection to all 50 million Ethiopians at risk for malaria through an ambitious plan to distribute long lasting insecticidal nets in all malarious areas by the end of 2007. The Carter Center was also asked to help in national efforts to monitor and evaluate the progress and effectiveness of the national control program. Since 2006, the program has built on TCC's existing programmatic networks in parts of Ethiopia while working closely with the MOH Malaria Control Program at national, regional, and local levels.

The Carter Center Focus on Malaria Control

The Carter Center's efforts in malaria control can be grouped into four focal areas:

1) Program Implementation and Interventions (treatment, net distribution and behavior change communications);
2) Monitoring and Evaluation

about stratification of data for analysis and targeting of interventions. The Carter Center has played an active role in the development of new standardized guidelines for epidemic monitoring in Ethiopia and

Annex 2: Neglected Tropical Diseases (Trachoma, Onchocerciasis and Lymphatic Filariasis)

TRACHOMA and its control

Trachoma is the world's leading infectious cause of preventable blindness. The World Health Organization estimates that 6 million people are blind due to trachoma, most of whom are women, and another 540 million are at risk of blindness or severe visual impairment. Blinding trachoma is caused by repeated infections of the conjunctiva (lining of the eye and eyelid) by the bacterium Chlamydia trachomatis.

Trachoma is transmitted from person to person through discharge from the ey -1.217 Td(an)2(d)2nose of infected individuals, which may be passed to others on hands, towels or clothing, or by fli -1.217 Tdwhich are attracted to ocular(an)2(d)2nasal discharge. Repeated infections lead to scarring of the conjunctiva which deforms the eyelid margin, causing ey -1.2lashes to turn inward and rub against the cornea. This condition, called trichiasis, causes severe pain and abrades the cornea, leading to other infections, opacity, and frequently to blindness.

Effective control of trachoma can be achieved using the SAFE strategy which consists of the following four components:

1) **S**urgery on eyelids to correct advanced stages of the disease;
2) **A**ntibiotics to treat active eye infection (predominantly oral azithromycin in the form of Zithromax®, donated by Pfizer Inc), which are given once per year during mass distributions;
3) **F**acial cleanliness, achieved through face washing, to prevent disease transmission; and
4) **E**nvironmental improvements to increase access to clean water and improved sanitation (building latrines to reduce fly br2.Tw -0.44 /P <(4)-3(r)111(1)-11(in)T13li ini/JTT1rJ[(d)N7a6tat7a6t as)5<</MCID 7

Annex 3: The Challenges of Disappearing Malaria: Lessons from the WHO Malaria Eradication Programme, 1955-1969

Presented by Dr. Randall Packard, Johns Hopkins University, Baltimore, MD

Remarkable successes in reducing the global burden of malaria have raised the possibility that malaria can be eliminated from many parts of the world. This optimism resembles that which led the WHO to launch the Global Programme for Malaria Eradication in the 1950s. While the Programme failed to achieve its goal, malaria was eliminated in a third of the countries in which it was attempted. Moreover, many other country programs came close to achieving elimination only to see malaria return. Both the

Annex 4:

Annex 5: MCP Publications and Abstracts since 2007

Publications

1) Endeshaw T, Graves PM, Ayele B, Mosher AW, Gebre T, Ayalew F, Genet A, Mesfin A, Shargie EB, Tadesse Z, Teferi T, Melak B, Richards RO and Emerson PM. (2012) Performance of Local Light Microscopy and the ParaScreen Pan/Pf Rapid Diagnostic Test to Detect Malaria in Health Centers in Northwest Ethiopia. PloS One 7(4): e33014. doi:10.1371/journal.pone.0033014.

2) Jima D, Wondabeku M, Alemu A, Teffera A, Awei N, Deressa W, Adissie A, Tadesse Z, Mosher AW, Richards FO and Graves PM. (2012) Analysis of malaria surveillance data in Ethiopia: what can be learned from the Integrated Disease Surveillance and Response System? Malaria Journal 11:330.

3) Graves PM, Ngondi JM, Hwang J, Getachew A, Gebre T, Mosher AW, Patterson AE, Shargie EB, Tadesse Z, Wolkon A, Reithinger R, Emerson PM, Richards FO Jr. (2011). Factors associated with mosquito net use by individuals in households owning nets in Ethiopia. Malaria Journal 10:354.

4) Ngondi JM, Graves PM, Gebre T, Mosher AW, Shargie EB, Emerson PM, and Richards FO. (2011). Which nets are being used: factors associated with mosquito net use in Amhara, Oromia andh(A)10(m)nSoM

5) Shargie EB, Ngondi J, Graves PM, Getachehw A, Hwang J, Gebr
Endeshaw T, Jima D, Tadesse Z, Tenaw E, Reithinger R, Emer
Ghebreyesus TA. (2010). Rapid increase
and decrease in prevalence of malaria in th

Annex 7: Malaria Program Review Meeting Agenda

4th Annual Malaria Program Review March 8, 2013		
8:00	Shuttle pickup at hotel	
8:30 – 9:00	Continental breakfast	
9:00 – 9:05 9:05 – 9:15	Morning session Welcome and introductions Overview	Dr. Paul Emerson (chair) Dr. Donald Hopkins Dr. Frank Richards
9:15 - 9:35 9:35 - 9:45	Ethiopia National Malaria Activities Report Discussion	Ms. Hiwot Solomon (FMOH)
9:45 - 10:15 10:15 -10:30	Carter Center Ethiopia	