

• ScienceDirect

# 1. Introduction

*Chlamydia trachomatis* 2.5% 1.5% 2.0% 1.0% 2.001 2.001 11 14 0.1% 3.5% 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

# 2. Materials and methods

## 2.1. The study population and sampling

2.001 2.001 11 14 5.0%

1.9 10% 9.5% 2.5% 15 1.5% 9.5% 2. (1) (2) (3) 2.000 22 1( 10





Figure 2






Table 4. Comparison of the prevalence of *S. pneumoniae* carriage in the nose and throat of children in the study population (n = 11155) and in the control population (n = 1444) in the 11155 study population.

Location	Study Population (n = 11155)	Control Population (n = 1444)	Prevalence (%)	95% CI	P-trend
Nose	4308	9	0.2	1.0	$P_{trend} < 0.001$
Throat	4490	9	1.1	0.5	4.1 1.4
Both	2342	9	3.3	1.5	0.4 34.4
Upper respiratory tract	555	4	1.2	1.0	
Lower respiratory tract	550	9	1.1	1.5	1.1 2.2
Upper respiratory tract (%)	351	44	1.3	1.0	$P_{trend} = 0.43$
Lower respiratory tract (%)	4130	93	1.5	1.0	
Upper respiratory tract (%)	150	2	1.1	1.3	
Lower respiratory tract (%)	9232	132	1.4	1.0	$P_{trend} = 0.49$
Upper respiratory tract (%)	154	29	1.4	0.4	1.3
Lower respiratory tract (%)	9	5	0.2	2.4	0.5 13.0
Upper respiratory tract (%)	208	39	1.4	1.0	$P_{trend} = 0.9$
Lower respiratory tract (%)	142	32	2.2	1.5	0.2
Upper respiratory tract (%)	490	95	1.4	1.0	0.1
Lower respiratory tract (%)	4031	41	1.0	1.0	$P_{trend} = 0.03$
Upper respiratory tract (%)	101	2	1.4	1.2	0.4 2.2
Lower respiratory tract (%)	5423	9	1.1	1.4	1.0 2.4
Upper respiratory tract (%)	55	0	0.1	1.0	$P_{trend} < 0.001$
Lower respiratory tract (%)	2102	44	3.0	2.5	1.4 3.9
Upper respiratory tract (%)	33	20	5.2	3.4	1.1 12.2
Lower respiratory tract (%)	112	12	1.0	1.1	2.5 1.9
Upper respiratory tract (%)	4	9	1.4	1.0	
Lower respiratory tract (%)	440	49	1.4	1.0	0.4 1.4
Upper respiratory tract (%)	114	102	1.4	1.0	0.93
Lower respiratory tract (%)	4041	44	1.4	1.2	0.350
Upper respiratory tract (%)	521	4	1.2	1.0	
Lower respiratory tract (%)	10434	140	1.5	0.4	0.2 2.1
Upper respiratory tract (%)	303	25	0.1	1.0	
Lower respiratory tract (%)	02	141	1.1	1.3	0.2 2.3
Upper respiratory tract (%)	1331	05	0.5	1.0	
Lower respiratory tract (%)	925	1	0.1	0.2	0.01 1.9
Upper respiratory tract (%)	943	1	0.1	0.2	0.02 1.9



A musical staff with a treble clef and a key signature of one flat. The notation includes various note values, rests, and dynamic markings. A 'fi' marking is present above the first measure. A '30' marking is located below the staff in the middle. The staff ends with a double bar line and repeat dots.





2004. doi: 10.1186/1475287520041414
31. <https://doi.org/10.1186/1475287520041414>
32. *Soc Sci Med* 1993; 37:1-5.
33. *Clin Microbiol Rev* 2004; 17:92-101.
34. *Lancet* 2008; 362:192-204.